

Please write clearly in	n block capitals.
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	I declare this is my own work.

# GCSE CHEMISTRY

H

Higher Tier Paper 2

Time allowed: 1 hour 45 minutes

#### **Materials**

For this paper you must have:

- a ruler
- · a scientific calculator
- the periodic table (enclosed).

## Instructions


- Use black ink or black ball-point pen.
- Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

### Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
TOTAL		








0 1

This question is about fuels and energy.

**Figure 1** shows the percentage of electricity generated in the UK between 2007 and 2017 using:

- oil
- · solar energy.

Figure 1



**0** 1. 1 Describe the changes in the percentage of electricity generated in the UK between 2007 and 2017 using:

- oil
- solar energy.

Use data from Figure 1 in your answer.

[3 marks]



0 1.2	Oil contains carbon and some sulfur.			
	When oil is burned, the products of combustion may be released into the atmosphere.			
	Explain the environmental effects of releasing these products of combustion into the atmosphere.			
	[6 marks]			



0 1.3	Suggest <b>one</b> reason why using solar energy is a more sustainable way of generating electricity than burning oil.  [1 mark]	outside box
0 1.4	Solar energy may <b>not</b> be able to replace the generation of electricity from fossil fuels completely.	
	Suggest <b>two</b> reasons why. [2 marks]	
	1	
	2	12
	Turn over for the next question	

0 2

This question is about alkanes.

**Table 1** shows information about some alkanes.


Table 1

Number of carbon atoms in alkane molecule	Boiling point of alkane in °C
4	0
5	36
6	69
7	х
8	126
9	151

0 2. 1 Plot the data from Table 1 on Figure 2.

[2 marks]

Figure 2





0 2 . 2	Predict the boiling point <b>X</b> of the alkane with seven carbon atoms in a molecu	ıle.
	Use <b>Table 1</b> and <b>Figure 2</b> .	
		[1 mark]
	<b>X</b> =°C	
0 2 . 3	Figure 2 is not suitable to show the boiling point of the alkane with three carb atoms in a molecule.	oon
	Suggest <b>one</b> reason why.	[1 mark]
0 2.4	What is the state at 20 °C of the alkane with four carbon atoms in a molecule	?
0 2.4	What is the state at 20 °C of the alkane with four carbon atoms in a molecule Use <b>Table 1</b> .	
0 2.4		? [1 mark]
0 2.4		
0 2 . 4		
0 2 . 4		
0 2 . 4		
0 2 . 4	Use Table 1.	
0 2 . 4		
0 2 . 4	Use Table 1.	
0 2 . 4	Use Table 1.	
0 2 . 4	Use Table 1.	
0 2 . 4	Use Table 1.	
0 2 . 4	Use Table 1.	



**Table 1** is repeated below.

Table 1

Number of carbon atoms in alkane molecule	Boiling point of alkane in °C
4	0
5	36
6	69
7	x
8	126
9	151

The alkane with nine carbon atoms in a molecule is called nonane.

0 2 . 5	Complete the formula of nonane.	[1 mark]
	C <sub>9</sub> H	
0 2.6	Nonane will condense lower in a fractionating column during fractional distitution the other alkanes in <b>Table 1</b> .	llation
	Explain why.	
	You should refer to the temperature gradient in the fractionating column.	[2 marks]



8

Do not write outside the Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 3	This question is about paper chromatography.
	A food colouring contains a dye.
0 3.1	Plan an investigation to determine the $R_{\mbox{\scriptsize f}}$ value for the dye in this food colouring.
	$R_f = \frac{\text{distance moved by substance}}{\text{distance moved by solvent}}$
	Your plan should include the use of:
	• a beaker
	• a solvent
	chromatography paper.
	[6 marks]



Do not write outside the box

0 3.2	Two students investigated a dye in a food colouring using paper chromatography.	out
	Each student did the investigation differently.	
	The R <sub>f</sub> values they determined for the <b>same</b> dye were different.	
	How did the students' investigations differ?	
	Tick (✓) one box.	
	Different length of paper used	
	Different period of time used	
	Different size of beaker used	
	Different solvent used	
0 3.3	Paper chromatography involves a stationary phase.	
	What is the stationary phase in paper chromatography?  [1 mark]	
	Tick (✓) one box.	
	Beaker	
	Dye	
	Paper	
	Solvent	



- This question is about poly(ethene) and polyesters.
- **0 4 . 1** Poly(ethene) is produced from ethene.

Figure 3 shows part of the displayed structural formula equation for the reaction.

Complete Figure 3.

[2 marks]

Figure 3


0 4 . 2	Poly(ethene) is a thermosoftening polymer.	
	Suggest why poly(ethene) is easier to recycle than thermosetting polymers.	[2 marks]
0 4.3	Ethene produces different forms of poly(ethene).	
	How can different forms of poly(ethene) be produced from ethene?	[1 mark]



- 0 4. Two different forms of poly(ethene) are:
  - high density poly(ethene) (HDPE)
  - low density poly(ethene) (LDPE).

Figure 4 represents part of the structures of HDPE and LDPE.

Figure 4



Explain why HDPE has a higher density than LDPE.	[2 marks]

Question 4 continues on the next page.



Figure 5 shows three monomers, A, B and C.

Monomer A can react with monomer B and with monomer C to produce polyesters.

Figure 5

0 4 . 5 Draw a circle on **Figure 5** around an alcohol functional group.

[1 mark]

- 0 4 . 6 Complete **Table 2** to show the formula of the small molecule produced when:
  - monomer A reacts with monomer B
  - monomer A reacts with monomer C.

[1 mark]

#### Table 2

Reacting monomers	Formula of small molecule produced
<b>A</b> and <b>B</b>	
A and C	

9



Do not write outside the Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 5	This question is about fertilisers.	
	Some fertilisers are described as NPK fertilisers because they contain three elements needed for healthy plant growth.	
0 5.1	Which <b>two</b> compounds each contain <b>two</b> of these elements?  [2 marks]	]
	Tick (✓) <b>two</b> boxes.	
	Ammonium nitrate	
	Ammonium phosphate	
	Calcium chloride	
	Calcium phosphate	
	Potassium chloride	
	Potassium nitrate	
0 5.2	Rocks containing calcium phosphate are treated with acid to produce soluble salts that can be used as fertilisers.	
	Name the soluble salts produced when calcium phosphate reacts with:	
	nitric acid	
	phosphoric acid.	
	[2 marks	]
	Nitric acid	_
	Phosphoric acid	



0 5.3	Ammonium sulfate is a compound in fertilisers.
	Ammonium sulfate can be made using an industrial process or in the laboratory.
	In the industrial process, the following steps are used.
	React streams of ammonia solution and sulfuric acid together.
	2. Evaporate the water by passing the solution down a warm column.
	3. Collect dry crystals continuously at the bottom of the column.
	In the laboratory, the following steps are used.
	1. React ammonia solution and sulfuric acid in a conical flask.
	2. Evaporate water from the solution until crystals start to form.
	3. Leave to cool and crystallise further.
	4. Separate the crystals using filtration.
	5. Dry the crystals between pieces of filter paper.
	Evaluate the two methods for producing a large mass of ammonium sulfate.  [4 marks]



Turn over ▶

8

0 6	This question is about cyclo	alkenes.	
	Cycloalkenes are ring-shap double carbon-carbon bond		s containing a
	Cycloalkenes react in a sim	ilar way to alkenes.	
0 6 . 1	Describe a test for the doub	le carbon-carbon bond in	cycloalkene molecules
		ile carbon-carbon bond in	cycloaikerie molecules.
	Give the result of the test.		[2 marks]
	Test		
	Result		
0   6  . 2	Table 3 shows the name ar	nd formula of three cycloa	lkenes.
	Table 3		
	Name	Formula	
	Cyclobutene	C <sub>4</sub> H <sub>6</sub>	
	Cyclopentene	C <sub>5</sub> H <sub>8</sub>	
	Cyclohexene	C <sub>6</sub> H <sub>10</sub>	
	Determine the general form	ula for cycloalkenes	
	Determine the general form	ula loi cycloaikelles.	[1 mark]
		Can aval farmania –	
		General formula = _	



Figure 6 shows the displayed structural formula of cyclohexene,  $C_6H_{10}$ 

## Figure 6

Chlorine reacts with cyclohexene to produce a compound with the formula C<sub>6</sub>H<sub>10</sub>Cl<sub>2</sub>

0 6. 3 Complete **Figure 7** to show the displayed structural formula of C<sub>6</sub>H<sub>10</sub>Cl<sub>2</sub>

[2 marks]

### Figure 7

$$egin{array}{cccc} \mathsf{H} & \mathsf{C} & \mathsf{C} \\ \mathsf{H} & \mathsf{C} & \mathsf{C} \\ \mathsf{H} & \mathsf{C} & \mathsf{C} \end{array}$$

0 6 . 4	Calculate the percentage by mass of chlorine in a molecule of C <sub>6</sub> H <sub>10</sub> Cl <sub>2</sub>				
	Relative atomic masses $(A_r)$ :	H = 1	C = 12	Cl = 35.5	[3

[3 marks]

Percentage by mass =	0/

8



0 7	Potash alum is a chemical compound.
	The formula of potash alum is KAl(SO <sub>4</sub> ) <sub>2</sub>
0 7.1	Give a test to identify the Group 1 metal ion in potash alum.
	You should include the result of the test.  [2 marks]
	Test
	Result
0 7.2	Name <b>one</b> instrumental method that could identify the Group 1 metal ion <b>and</b> show
	the concentration of the ion in a solution of potash alum.  [1 mark]

	A student identifies the other metal ion in potash alum.	outside box
	The student tests a solution of potash alum by adding sodium hydroxide solution until a change is seen.	
0 7.3	Give the result of this test.  [1 mark]	
0 7.4	This test gives the same result for several metal ions.	
	What additional step is needed so that the other metal ion in potash alum can be identified?	
	Give the result of this additional step.  [2 marks]	
	Additional step	
	Result	
0 7.5	Describe a test to identify the presence of sulfate ions in a solution of potash alum.	
	Give the result of the test.  [3 marks]	
	Test	
	Result	
		9



0	8	This guestion is about copper and alloys of copper
U	O	This question is about copper and alloys of copper

Solders are alloys used to join metals together.

Some solders contain copper.

Table 4 shows information about three solders, A, B and C.

Table 4

Solder	Melting point in °C	Metals in solder
Α	183	tin, copper, lead
В	228	tin, copper, silver
С	217	tin, copper, silver

0 8.1	Solder <b>B</b> and solder <b>C</b> are now used more frequently than solder <b>A</b> for health	n reasons.
	Suggest <b>one</b> reason why.	
	Use <b>Table 4</b> .	[1 mark]
		[ i iliai k]
0 8.2	Suggest <b>one</b> reason why solders <b>B</b> and <b>C</b> have different melting points.	
	Use <b>Table 4</b> .	[1 mark]



Do	not	V	vr	ite
ou	tside	9	tr	ie
	ho	¥		

	Copper can be obtained by:
	processing copper ores
	recycling scrap copper.
0 8.3	Suggest <b>three</b> reasons why recycling scrap copper is a more sustainable way of obtaining copper than processing copper ores.
	[3 marks]
	1
	2
	3

Question 8 continues on the next page



	Copper is extracted from low-grade ores by phytomining.		outsi b
0 8.4	Describe how copper is extracted from low-grade ores by phytomining.	[4 marks]	
0 8.5	Phytomining has <b>not</b> been widely used to extract copper.		
	Suggest <b>two</b> reasons why.	[2 marks]	
	1		
	2		
			1



0 9

A student investigated how a change in concentration affects the rate of the reaction between zinc powder and sulfuric acid.

The equation for the reaction is:


$$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + H_2(g)$$

This is the method used.

- 1. Pour 50 cm<sup>3</sup> of sulfuric acid of concentration 0.05 mol/dm<sup>3</sup> into a conical flask.
- 2. Add 0.2 g of zinc powder to the conical flask.
- 3. Put the stopper in the conical flask.
- 4. Measure the volume of gas collected every 30 seconds for 5 minutes.
- 5. Repeat steps 1 to 4 with sulfuric acid of concentration 0.10 mol/dm3

Figure 8 shows the apparatus used.

Figure 8



0 9.

The student made an error in setting up the apparatus in Figure 8.

What error did the student make?


[1 mark]



The student corrected the error.

Figure 9 shows the student's results.

Figure 9



0 9 . 2	Explain why the lines of best fit on <b>Figure 9</b> become horizontal.	[2 marks]

0 9.3 How does **Figure 9** show that zinc powder reacts more slowly with 0.05 mol/dm³ sulfuric acid than with 0.10 mol/dm³ sulfuric acid? [1 mark]



Do not write outside the box

0 9 . 4	Determine the rate of the reaction for 0.05 mol/dm³ sulfuric acid at 80 seconds.	OL
	Show your working on <b>Figure 9</b> .	
	Give your answer to 2 significant figures.	
	[5 marks]	
	Rate of reaction (2 significant figures) = cm³/s	
	( ° ° ° , s.ii , e	
0 9 . 5	The activation energy for the reaction between zinc and sulfuric acid is lowered if a	
0 0 .	solution containing metal ions is added.	
	What is the most likely formula of the metal ions added?  [1 mark]	
	Tick (✓) one box.	
	Al <sup>3+</sup>	
	Ca <sup>2+</sup>	
	Q 2+	
	Cu <sup>2</sup> '	
	Na <sup>+</sup>	-
	Cu <sup>2+</sup> Na <sup>+</sup>	



1 0	This question is about alkenes and alcohols.
	Ethene is an alkene produced from large hydrocarbon molecules.
	Large hydrocarbon molecules are obtained from crude oil by fractional distillation.
10.1	Name the process used to produce ethene from large hydrocarbon molecules.  [1 mark]
1 0 . 2	Describe the conditions used to produce ethene from large hydrocarbon molecules.  [2 marks]



1 0 . 3	Ethanol can be produced from ethene and steam.
	The equation for the reaction is:
	$C_2H_4(g) + H_2O(g) \rightleftharpoons C_2H_5OH(g)$
	The forward reaction is exothermic.
	Explain how the conditions for this reaction should be chosen to produce ethanol as economically as possible.  [6 marks]




Do not write outside the box

1 0.4	Ethanol can also be produced from sugar solution by adding yeast.	
	Name this process.	
	[1 n	nark]
1 0.5	Butanol can be produced from sugar solution by adding bacteria.	
	Sugar solution is broken down in similar ways by bacteria and by yeast.	
	Suggest the reaction conditions needed to produce butanol from sugar solution	
	by adding bacteria.	arks]
	įz mi	aiksj



	Ethanol and butanol can be used as fuels for cars.		
1 0.6	A car needs an average of 1.95 kJ of energy to travel 1 m		
	Ethanol has an energy content of 1300 kilojoules per mole (kJ/mol).		
	Calculate the number of moles of ethanol needed by the car to travel 200 km	n [3 marks]	
	Number of moles =	mol	
1 0.7	When butanol is burned in a car engine, complete combustion takes place.		
	Write a balanced equation for the complete combustion of butanol.		
	You do <b>not</b> need to include state symbols.	[2 marks]	
	END OF QUESTIONS		







Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



outside the There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED Copyright information For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk. Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful

and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.



Copyright © 2021 AQA and its licensors. All rights reserved.



Do not write