Please check the examination detail	ils bel	ow before ente	ring your candidate information
Candidate surname			Other names
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	Cen	tre Number	Candidate Number
Time 1 hour 45 minutes		Paper reference	1CH0/2F
Chemistry PAPER 2 Foundation Tier			
You must have: Calculator, ruler			Total Marks

Instructions

- Use **black** ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- In questions marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ▶







Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

1 (a) Figure 1 shows a mug made of clay ceramic.




Figure 1

Which property of clay ceramic makes it suitable for use as a mug?

(1)

- A is brittle
- **B** is not transparent
- **D** does not dissolve in water

(b) Figure 2 shows the properties of some materials.

	property					
material	brittle	conducts electricity	conducts heat	transparent		
clay ceramic	yes	no	no	no		
glass	yes	no	no	yes		
metal	no	yes	yes	no		
polymer	no	no	no	no		

Figure 2

Figure 3 shows an electrical cable.

The electrical cable is made of metal wire coated with another material.

The metal wire inside the electrical cable conducts electricity.




Figure 3

(i) Name a material from Figure 2 that would be suitable for coating the metal wire.

1)

(ii) Which type of particle moves through the metal wire to allow it to conduct electricity?

(1)

- **A** atoms
- **B** electrons
- C neutrons
- **D** protons

- (c) There are some concerns that nanoparticles may cause harm if they enter the human body.
 - (i) Suggest one way that nanoparticles can enter the human body.

(1)

(ii) Suggest one possible risk if nanoparticles enter the human body.

(1)

(iii) The surface area of a nanoparticle of gold is 150 nm². The volume of a nanoparticle of gold is 125 nm³.

$$ratio = \frac{surface area}{volume}$$

Calculate the surface area to volume ratio of this nanoparticle of gold.

(1)

surface area to volume ratio = :1

(Total for Question 1 = 6 marks)

BLANK PAGE



2	Magnesium reacts with dilute sulfuric acid to form magnesium sulfate and hydrogen	gas.
	A student wants to find out the effect of temperature on the rate of this reaction.	
	The student used the following method.	
	 step 1 pour 25 cm³ of dilute sulfuric acid into a conical flask step 2 warm the acid until its temperature is 30 °C step 3 add a piece of magnesium to the acid step 4 start a stopwatch step 5 wait until the reaction has finished step 6 stop the stopwatch step 7 repeat steps 1–6 but at 50 °C. (a) The student kept the volume of sulfuric acid the same when they repeated the 	
	method at 50°C.	
	State two other variables that should be kept the same.	(2)
1		,
1 2		
∠	(b) Which piece of equipment can be used to find the volume of 25 cm ³ of sulfuric ac	id?
		(1)
	■ A balance	
	■ B measuring cylinder	
	■ D thermometer	
	(c) State how the student will know that the reaction has finished.	(1)
	(d) The reaction at 50 °C was faster than the reaction at 30 °C.	
	Give one reason, in terms of particles, why the reaction at 50° C was faster than the reaction at 30° C.	
	the reaction at 50°C.	(1)



(Total for Question	on 2 = 7 marks)
average rate of reaction =	cm ³ s ⁻¹
	(2)
Calculate the average rate of reaction, in cm ³ s ⁻¹ , for the first 60 seconds reaction.	s of the
(e) At 50°C, 15.0 cm³ of gas was produced during the first 60 seconds of the	ne reaction.

- **3** This question is about the noble gases.
 - (a) (i) State, in terms of outer shell electrons, why the noble gases are unreactive.

(1)

(ii) Figure 4 shows an airship, filled with helium, floating above the ground.




Figure 4

Helium, hydrogen and krypton are gases. Figure 5 shows the reactivity and density, at room temperature and pressure, of helium, hydrogen and krypton.

gas	reactivity	density in g cm ⁻³
helium	unreactive	0.00018
hydrogen	very reactive	0.00009
krypton	unreactive	0.00380

Figure 5

The density of air is $0.001225 \,\mathrm{g\,cm^{-3}}$.

Helium is used in airships.

Explain why hydrogen and why krypton are **not** used in airships.

(3)

(b) Mendeleev produced one of the earliest periodic tables.

State why he could **not** include any of the noble gases in his periodic table.

(1)


(c) Figure 6 shows the boiling points of some of the noble gases.

noble gas	boiling point in °C
helium	-269
neon	-246
argon	-186
krypton	-153
xenon	

Figure 6

(i) Complete the bar chart to show the boiling points of neon and argon.

(2)



(ii) Predict the boiling point of xenon.

(1)

boiling point of xenon =°C

(Total for Question 3 = 8 marks)



- **4** A student carried out some tests on copper sulfate.
 - (a) A flame test was carried out on some copper sulfate crystals.

The student used the following method.

- **step 1** dip a wire in hydrochloric acid, then hold the wire in a roaring Bunsen burner flame
- **step 2** dip the wire in hydrochloric acid again, then dip the wire in the copper sulfate crystals
- **step 3** hold the wire with the copper sulfate in the roaring Bunsen burner flame.
- (i) State why in step 1 the wire is dipped in hydrochloric acid and held in a roaring Bunsen burner flame.

(1)

(ii) State why in step 2 the wire is dipped in hydrochloric acid again before dipping it in the copper sulfate crystals.

(1)

(iii) What colour should be seen when the flame test is carried out on copper sulfate?

(1)

- A blue-green
- **B** lilac
- C orange-red
- **D** yellow
- (b) A solution of the copper sulfate was tested in a flame photometer.
 - (i) Give an advantage of using a flame photometer, rather than a flame test, to test for copper ions.

(1)



(ii) Figure 7 shows some values obtained when producing a calibration graph for copper ions in a flame photometer.

Draw a line of best fit through the points.

(1)

display reading




Figure 7

(iii) A solution containing copper ions gave a display reading of 0.030.

Use the calibration graph in Figure 7 to find the concentration of copper ions in this solution.

(1)

concentration of copper ions = _____ppm

- (c) The student tested an unknown solution for sulfate ions by
 - 1. adding a few drops of dilute sulfuric acid to the unknown solution
 - 2. then adding a few drops of barium chloride solution.

A white precipitate was formed.

The student cannot conclude from this result that the unknown solution contains sulfate ions.

Explain what the student should change to obtain a valid result.

(2)

(d) Calculate the percentage by mass of copper in copper sulfate, CuSO₄.

(relative atomic masses: Cu = 63.5 relative formula mass of $CuSO_4 = 159.5$)

Give your answer to the nearest whole number.

(3)

percentage by mass of copper =

(Total for Question 4 = 11 marks)

BLANK PAGE




5					
	(a) (I)	State	tne	name given to the group 7 elements.	(1)
				ne other element that is in group 7. Deriodic table on the back of this exam paper to help you.	(1)
	(iii)	Whic	h ele	ement is liquid at room temperature and pressure?	(1)
		×	A	fluorine	
		×	В	chlorine	
		X	c	bromine	
		×	D	iodine	
	(iv)	Whic	h ele	ement is dark-grey in colour at room temperature and pressure?	(1)
		\times	A	fluorine	
		\times	В	chlorine	
		\times	C	bromine	
		X	D	iodine	
	` '			th chlorine to form tin chloride. tin chloride contains 1.19g of tin and 1.42g of chlorine.	
	Calc	culate	the	e empirical formula of this tin chloride.	
	(rela	ative a	ator	mic masses: Cl = 35.5, Sn = 119.0)	
	You	must	t sho	ow your working.	(3)
					(3)
				empirical formula of this tin chloride =	



(c) Tin also reacts with fluorine.

The reaction between fluorine and tin is much more vigorous than the reaction between chlorine and tin.

Figure 8 shows the electronic configurations of fluorine and chlorine.






Figure 8

Explain, in terms of their electronic configurations, why fluorine reacts with tin more vigorously than chlorine reacts with tin.

_	-	

(2)

(Total for Question 5 = 9 marks)



BLANK PAGE

6 Figure 9 shows a sample of hydrogen peroxide solution decomposing to form water and oxygen gas.




Figure 9

(a) (i) Write the word equation for hydrogen peroxide solution decomposing.

(1)


(ii) In this reaction hydrogen peroxide is a solution, water is a liquid and oxygen is a gas.

Draw one straight line from each substance to its correct state symbol.

(2)

substance

hydrogen peroxide solution






liquid water



(g)



(l)

(b) Describe the test to show the gas produced is oxygen.

(2)



(2)

(c) Figure 10 shows the electron arrangement for an atom of hydrogen and an atom of oxygen.




Figure 10

Complete the dot and cross diagram in Figure 11 for a molecule of water, H_2O . Draw outer shell electrons only.




Figure 11

(d) Liver contains the enzyme catalase.

A piece of liver was added to another sample of hydrogen peroxide solution.

Figure 12 shows the results.





Figure 12





- **7** This question is about alcohols.
 - (a) (i) Figure 14 shows an incomplete diagram of the structure of a molecule of propanol, CH₃CH₂CH₂OH.

Complete the structure of the molecule of propanol in Figure 14.

(2)




Figure 14

(ii) Propanol can be converted into propanoic acid.

What happens to propanol in this reaction?

(1)

- **B** it is neutralised
- C it is oxidised
- **D** it is polymerised

(b) A student used an alcohol burner to find the mass of different alcohols needed to raise the temperature of 100 cm³ of water by 20 °C.

Figure 15 shows their results.

alcohol	initial mass of alcohol burner and alcohol in g	final mass of alcohol burner and alcohol in g	mass of alcohol used in g
ethanol	122.51	122.02	0.49
propanol	168.55	168.13	0.42
butanol	152.62	152.23	
pentanol	67.22	66.86	0.36

Figure 15

(i) Calculate the mass of butanol used.

(1)

mass of butanol =g

*(ii) Figure 16 shows equipment that can be used to obtain the results shown in Figure 15.




Figure 16

Describe an experiment, using the equipment in Figure 16, that could be used to obtain results like those shown in Figure 15.

.....

(6)



				۲		
				1		
				1		
	G			n		
	А	Б		5		
	И		ù	а		
		S				
	ũ		7	٠		
	q	D				
	7					۰
	Κ.	_	۷	_	2	
	2	₹	z			
	Ì	d		2		
	Δ	-	۰	-	,	۰
	D				ú	
	ч	r			3	
	2	6	ú	ú		
		2	7	7		
	4	à	4	è		
	0	₹	7	₹		
					٩	
		١,		S	á	
	١		E	7	•	
		3				
	λ			Σ		
		7	7	7	,	
	Λ	'n	=	'n	ı	
		z	Į	7		
	9	9	٣	9	ę	
	à	ú	ė	ú		
		7				
	á	è	×	à		
	☇	2	∠	S		
>	S	ř	ź	Ř	į	
>	1	Ì	1	Ì	į	
	Į	Ì	1	ļ	ĺ	֡
	3		4	ļ		֡
	3		4	ļ		
			1	ļ		
			1	ļ		
				ļ		
				ļ		
				ļ		
				ļ		
	3	2		ļ		
	3	2		ļ		
		2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
	3	2		ļ		
				ļ		
				ļ		
	3			ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		
				ļ		



(iii) The results show that $0.36\mathrm{g}$ of pentanol was needed to raise the temperature of the water by $20^\circ\mathrm{C}$.	ıre
Calculate the mass of pentanol needed to raise the temperature of water by	y 1°C.
Give your answer to 2 decimal places.	
Show your working.	(2)
	(2)
mass of pentanol =	g
(Total for Question 7 = 12	marks)

8 (a) The concentration of a solution can be calculated using the equation

concentration of solution =
$$\frac{\text{mass of solid}}{\text{volume of solution}}$$


A student dissolved 9.25 g of ammonium chloride in water and made up the solution to a volume of 200 cm³.

Use the equation to calculate the concentration of this solution in g dm⁻³.

(2)

concentration =g dm⁻³

(b) Dissolving ammonium chloride in water is an endothermic process. Figure 17 shows part of the reaction profile for this process.



progress of reaction

Figure 17

(i) Explain how Figure 17 shows that dissolving ammonium chloride in water is an endothermic process.

(2)


 	 ••••••	 ••••••	 	 	 ••••••	 	 •••••	



(ii) Complete the reaction profile in Figure 17 and label the activation energy.

(2)

(c) A student used the equipment in Figure 18 to investigate whether electricity can pass through solid ammonium chloride and through ammonium chloride solution.






Figure 18

If an electrical current flows in the circuit, the lamp will light up.

Figure 19 shows the results of the investigation.

substance	lamp
solid ammonium chloride	did not light up
ammonium chloride solution	lit up brightly

Figure 19

(3)

- (d) Ammonia gas is toxic.
 - (i) Which symbol should be placed on a container of a toxic gas?

(1)









(ii) Give **one** safety precaution that should be taken when working with toxic gases in the laboratory.

(1)

(Total for Question 8 = 11 marks)

/ _ \			a mixture of hydrocarbons that can be obtained from crude oil.	
(a)	Stat	te th	e name of the process used to separate diesel oil from crude oil.	(1)
(b)			il contains alkanes. Ikanes are part of an homologous series.	
	Whi	ich s	tatement about compounds in this homologous series is true?	(4)
	X	Α	they have the same chemical formula	(1)
	X	В	they have the same empirical formula	
	X	C	they have the same general formula	
	X	D	they have the same molecular formula	
(c)			uels such as diesel oil are burned, the high temperatures produced can itrogen and oxygen in the air to form the pollutant nitrogen dioxide.	
	Con	nple	te the balanced equation for the reaction.	(2)
			$N_2 + 2O_2 \rightarrow \dots$	
(d)	-		how the greenhouse effect is caused by the gases produced by the te combustion of diesel oil.	
		•		(3)

*(e)	Diesel oil can contain impurities of sulfur. Burning diesel oil containing impurities of sulfur can result in acid rain. Acid rain is harmful to the environment.	
	Explain how acid rain is formed and the harm it can do.	(6)



10 This question is about polymers.

(a) (i) State a problem with **recycling** polymers.

(1)

(ii) Describe a problem associated with the **disposal** of polymers.

(2)

(b) Poly(chloroethene) is a polymer made from chloroethene. A molecule of chloroethene is shown in Figure 20.

Figure 20

(i) On Figure 20, draw a circle around the functional group in this molecule.

(1)

(ii) Draw a section of a poly(chloroethene) molecule containing three repeating units, showing all bonds.

(3)

(iii) What type of polymer is poly(chloroethene)?	
	(1)
(iv) Calculate the relative formula mass of a poly(chloroethene) molecule m from 2850 chloroethene molecules, C ₂ H ₃ Cl.	ade
(relative atomic masses: H = 1.00, C = 12.0, Cl = 35.5)	
Give your answer to three significant figures.	
Show your working.	(0)
	(3)
relative formula mass =	
(Total for Question 10 =	: 11 marks)
TOTAL FOR PAPER = 1	00 MARKS



The periodic table of the elements

0 4 He helium 2	20 Ne neon 10	40 Ar argon 18	84 Kr ypton 36	131 Xe xenon 54	[222] Rn radon 86
7	19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85
9	16 O 0xygen 8	32 S sulfur 16	79 Selenium 34	128 Te tellurium 52	[209] Po polonium 84
ιΩ	14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83
4	12 C carbon 6	28 Si silicon 14	73 Ge germanium 32	119 Sn tin 50	207 Pb lead 82
ო	11 B boron 5	27 AI aluminium 13	70 Ga gallium 31	115 In indium 49	204 T thallium 81
·			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80
			63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79
			59 nickel 28	106 Pd palladium 46	195 Pt platinum 78
			59 Co cobalt 27	103 Rh rhodium 45	192 Ir iridium 77
1 Hydrogen			56 iron 26	101 Ru ruthenium 44	190 Os osmium 76
			55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75
	nass ool umber	(protor) manber	52 Cr	96 Mo molybdenum 42	184 W tungsten 74
Key	relative atomic mass atomic symbol name atomic (proton) number		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73
	relativ ato atomic		48 Ti titanium 22	91 Zr zirconium 40	178 Hf hafnium 72
·			45 Sc scandium 21	89 × yttrium 39	139 La* lanthanum 57
2	9 Be beryllium 4	24 Mg magnesium 12	40 Ca calcium 20	88 Sr strontium 38	137 Ba barium 56
-	7 Li lithium 3	23 Na sodium 11	39 potassium 19	85 Rb rubidium 37	133 Cs caesium 55

^{*} The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.